

**SANTA BARBARA SURGERY CENTER
OPHTHALMOLOGY ADMISSION ORDERS
FAX: 866-297-5257**

PLEASE PRINT LEGIBLY

PATIENT NAME _____ ADMISSION DATE: _____

ADMITTING MD: _____

- SURGICAL CONSENT:**
- Cataract Extracton _____ eye with possible Intraocular Lens Implant
 - Clear Lensectomy _____ eye
 - Limbal Relaxing Incision _____ eye
 - Other _____ eye

History & Physical Dictated BY: Surgeon Internist

LABORATORY STUDIES:

- No ECG Preoperative EKG ordered Where? _____
- Obtain Blood Glucose for Diabetic Patients Hemacue (HGB) Urine Pregnancy Test
- Diagnosis Related Labs Requested: Renal Panel Hb/Hct PT/PTT
- Lab has been ordered Where? _____

PREOPERATIVE ORDERS:

- Obtain ECG (If not done within 90 days) for patients > 45 except Cataracts and Refractive Lensectomy
- Betadine face prep prior to surgery (if patient allergic, use Hibiciens)
- Prior to betadine prep or dilation drops, instill one drop proparacaine ophthalmic solution in operative eye.
- Instill each checked ophthalmic solution drops into the patient's operative eye every 5-10 minutes X _____
May discontinue when patient is well dilated.
- Cyclogyl 2% Neosynephrine 2.5% Pilocarpine 2%
- Mydracil 1% Vigamox 0.5% Mydracil 0.33% / Cyclogyl 0.66% / Neosynephrine 3.33%
- Neosynephrine 10% Zymar 0.3% _____

2% Lidocaine Jelly. 2 times preop TOPICAL

Other medications: _____

Physician's Routine Block Nurse to draw up block Physician to draw block

- Lidocaine 2% with Epi 1:1000,000 = 5 ml
- 0.75% Marcaine Plain PF 0.5 ml + 2% Lidocaine Plain 0.5 ml = 1 ml
- Lidocaine 2% Plain (2.5 ml) + 0.75% Marcaine (2.5 ml) = 5 ml
- Lidocaine 2% Plain (5 ml) + 0.75% Marcaine (5 ml) = 10 ml
- 10 ml BSS w/ 0.2 ml epi 1:1000 PF (0.5 ml) + 1% Lidocaine PF (0.5 ml) = 1 ml
- Mepivacaine 1.5% (9 ml) + Sodium Bicarb 8.4% (1 ml) = 10 ml
- Mepivacaine 1.5% (4 ml) + Bupivacaine 0.75% (4 ml) + Vitrase 100 units (0.5 ml) = 8.5 ml

Follow routine anesthesia orders

Other: _____

SIGNATURE: _____ MD TIME: _____ DATE: _____

